



## Review Article

## Cannabidiol as an emergent therapeutic strategy for lessening the impact of inflammation on oxidative stress

George W. Booz\*

Department of Pharmacology and Toxicology, School of Medicine, and Center for Excellence in Cardiovascular–Renal Research, University of Mississippi Medical Center, Jackson, MS 39216, USA

## ARTICLE INFO

## Article history:

Received 9 September 2010

Revised 4 January 2011

Accepted 5 January 2011

Available online 14 January 2011

## Keywords:

Inflammation

Oxidative stress

Immune system

Metabolic syndrome

Endocannabinoid

Free radicals

## ABSTRACT

Oxidative stress with reactive oxygen species generation is a key weapon in the arsenal of the immune system for fighting invading pathogens and initiating tissue repair. If excessive or unresolved, however, immune-related oxidative stress can initiate further increasing levels of oxidative stress that cause organ damage and dysfunction. Targeting oxidative stress in various diseases therapeutically has proven more problematic than first anticipated given the complexities and perversity of both the underlying disease and the immune response. However, growing evidence suggests that the endocannabinoid system, which includes the CB<sub>1</sub> and CB<sub>2</sub> G-protein-coupled receptors and their endogenous lipid ligands, may be an area that is ripe for therapeutic exploitation. In this context, the related nonpsychotropic cannabinoid cannabidiol, which may interact with the endocannabinoid system but has actions that are distinct, offers promise as a prototype for anti-inflammatory drug development. This review discusses recent studies suggesting that cannabidiol may have utility in treating a number of human diseases and disorders now known to involve activation of the immune system and associated oxidative stress, as a contributor to their etiology and progression. These include rheumatoid arthritis, types 1 and 2 diabetes, atherosclerosis, Alzheimer disease, hypertension, the metabolic syndrome, ischemia–reperfusion injury, depression, and neuropathic pain.

© 2011 Elsevier Inc. All rights reserved.

## Contents

Introduction . . . . .	1055
Biochemistry of cannabidiol . . . . .	1055
Mechanisms of action . . . . .	1055
Actions on immune cells . . . . .	1056
Pain . . . . .	1056
Diabetes and diabetic complications . . . . .	1056
Hypertension . . . . .	1056
Ischemia–reperfusion injury . . . . .	1057
Depression . . . . .	1057
Neurodegenerative diseases . . . . .	1057
Obesity and the metabolic syndrome . . . . .	1057
Atherosclerosis . . . . .	1058
Conclusions . . . . .	1059
Acknowledgments . . . . .	1059
References . . . . .	1059

**Abbreviations:** 5-HT<sub>1A</sub> receptor, 5-hydroxytryptamine (serotonin) receptor subtype 1A; A<sub>2A</sub>, adenosine A<sub>2A</sub> receptor (ADORA2A); Abn-CBD, abnormal-cannabidiol; Aβ, β-amyloid peptide; BHT, butylated hydroxytoluene; CB<sub>1</sub>, cannabinoid receptor type 1; CB<sub>2</sub>, cannabinoid receptor type 2; CBD, cannabidiol; CD36, cluster of differentiation 36; FAAH, fatty acid amide hydrolase; ICAM-1, intercellular adhesion molecule 1; IL, interleukin; iNOS (or NOS2), inducible NOS; LDL, low-density lipoprotein; LOX, lipoxygenase; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; MCP1, monocyte chemoattractant protein 1; mmLDL, minimally modified LDL; NF-κB, nuclear factor κ-light-chain enhancer of activated B cells; NK, natural killer; NKT, natural killer T; NOD mouse, nonobese diabetic mouse; oxLDL, oxidized LDL; PPAR, peroxisome proliferator-activated receptor; PVN, paraventricular nucleus; Rap1, Ras-related protein 1; ROS, reactive oxygen species; STAT, single transducer and activator of transcription; Tc, cytotoxic T; Th, T helper; TNFα, tumor necrosis factor-α; VCAM-1, vascular cell adhesion molecule 1; Δ<sup>9</sup>-THC, Δ<sup>9</sup>-tetrahydrocannabinol.

\* Fax: +1 601 984 1637.

E-mail address: [gbooz@umc.edu](mailto:gbooz@umc.edu).

## Introduction

(–)-Cannabidiol (CBD) is the major nonpsychotropic cannabinoid compound derived from the plant *Cannabis sativa*, commonly known as marijuana. CBD was first isolated in 1940 and its structure and stereochemistry were determined in 1963 [1,2]. Interest in exploiting CBD therapeutically was initially focused on its interactions with the primary psychotropic ingredient of *Cannabis*,  $\Delta^9$ -THC, and its sedative and antiepileptic effects and later its antipsychotic and anxiolytic actions and utility in treating movement disorders [3]. As chronicled elsewhere [3], the past several years have seen a renewed interest in CBD because of the discovery of its antioxidative, anti-inflammatory, and neuroprotective effects, actions that occur for the most part independent of the canonical cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptors [1,4]. CBD may prove to have therapeutic utility in a number of conditions involving both inflammation and oxidative stress, including Parkinson disease, diabetes, rheumatoid arthritis, Alzheimer disease, and ischemia-reperfusion injury.

The contribution of the endocannabinoid system to inflammation and regulation of the immune system is an area of intense study that is beyond the scope of this article, and the reader is referred to several recent excellent reviews [5–8]. However, a brief overview of the system is helpful in discussing CBD. The endocannabinoid system comprises the following: (1) the G-protein-coupled cannabinoid receptors CB<sub>1</sub> and CB<sub>2</sub>, which are located in both the central nervous system and the periphery; (2) their arachidonate-based lipid ligands, e.g., 2-arachidonoylglycerol and anandamide (*N*-arachidonylethanolamine); and (3) the enzymes that synthesize and degrade these ligands. The endocannabinoid system plays a role in a variety of physiological processes including appetite, pain sensation, and mood. Evidence indicates that both CB<sub>1</sub> and CB<sub>2</sub> are expressed by cells of the immune system and are upregulated in the activation state. Levels of CB<sub>2</sub> appear to be higher than those of CB<sub>1</sub>, with decreasing amounts of CB<sub>2</sub> in human B cells, natural killer (NK) cells, monocytes, polymorphonuclear neutrophils, and T cells [6]. Macrophages and related cells, microglia and osteoclasts, express both cannabinoid receptors. CB<sub>2</sub> activation of immune cells is associated with changes in cytokine release and migration [6].

## Biochemistry of cannabidiol

CBD (Fig. 1) is a resorcinol-based compound that was shown to have direct, potent antioxidant properties by cyclic voltammetry and a spectrophotometric assay of oxidation in a Fenton reaction [9]. In an *in vitro* glutamate neuronal toxicity model, CBD was shown to be more protective than either  $\alpha$ -tocopherol or vitamin C and comparable to butylated hydroxytoluene (BHT); although as noted by the authors, CBD, unlike BHT, does not seem to promote tumors [9]. CBD was also reported to act as an antioxidant at submicromolar concentrations in preventing serum-deprived cell death of cultured human B lymphoblastoid and mouse fibroblast cells [10]. The antioxidant chemistry of CBD may have utility *in vivo* as well. The




Fig. 1. Chemical structure of cannabidiol (CBD).

protective effects of CBD in a rat binge ethanol-induced brain injury model [11] and a rat model of Parkinson disease [12] were ascribed to its antioxidant properties. As will become clear from this review, however, the antioxidant actions ascribed to CBD in various *in vivo* models of human diseases probably exceed those attributable to its chemistry alone. Rather, the therapeutic antioxidant properties of CBD would seem to result in no small measure from its modulation of cell signaling events that underlie the self-sustaining cycle of inflammation and oxidative stress.

## Mechanisms of action

Several interactions with relevance to the immune system and oxidative stress are discussed here. First, despite having low affinity for CB<sub>1</sub> and CB<sub>2</sub> receptors, CBD has been shown to antagonize the actions of cannabinoid CB<sub>1</sub>/CB<sub>2</sub> receptor agonists in the low-nanomolar range, consistent with noncompetitive inhibition [13]. At 1–10  $\mu$ M, CBD appears to function as an inverse agonist at both CB<sub>1</sub> and CB<sub>2</sub> receptors [13]. Second, CBD acts as an inhibitor (IC<sub>50</sub> 28  $\mu$ M) of fatty acid amide hydrolase (FAAH), the major enzyme for endocannabinoid breakdown. Because FAAH activity correlates with gastrointestinal mobility, CBD may have utility in treating intestinal hypermotility associated with certain inflammatory diseases of the bowel [14].

Third, CBD is a competitive inhibitor, with an IC<sub>50</sub> in the nanomolar range, of adenosine uptake by the equilibrative nucleoside transporter 1 of macrophages and microglial cells, the resident macrophage-like immune cells of the brain. By increasing exogenous adenosine, which in turn activates the A<sub>2A</sub> adenosine receptor, CBD exerts immunosuppressive actions on macrophages and microglial cells as evidenced by decreased tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) production after treatment with lipopolysaccharide (LPS) [15,16]. CBD may thus be of benefit in treating neurodegenerative diseases associated with hyperactivation of microglial, as well as retinal neuroinflammation seen in such conditions as uveitis, diabetic retinopathy, age-related macular degeneration, and glaucoma. Note, however, that adenosine activates other receptors in addition to A<sub>2A</sub> that often have opposing consequences on immune regulation and inflammation [17,18]. In several *in vivo* models of neurodegeneration or inflammation, moreover, the beneficial effects of CBD were demonstrated not to involve adenosine receptors.

Fourth, CBD has been shown to have potent actions in attenuating oxidative and nitrosative stress in several human disease models, although the exact mechanism is unclear. For instance, CBD pretreatment was found to attenuate high-glucose-induced mitochondrial superoxide generation and NF- $\kappa$ B activation in human coronary artery endothelial cells, along with nitrotyrosine formation and expression of inducible nitric oxide synthase (iNOS) and adhesion molecules ICAM-1 and VCAM-1 [19]. Notably, high-glucose-induced transendothelial migration of monocytes, monocyte-endothelial adhesion, and barrier disruption were attenuated as well. These findings lend support to the conclusion that that CBD may have therapeutic utility in treating diabetic complications and atherosclerosis. In another study, CBD was reported to reduce expression of reactive oxygen species (ROS)-generating NADPH oxidases, as well as iNOS and nitrotyrosine generation, in a cisplatin nephropathy model *in vivo*, consequently lessening cell death in the kidney and improving renal function [20]. From these studies, it is tempting to speculate that CBD may act directly at the level of the mitochondrion or nucleus to oppose oxidative/nitrosative stress.

Fifth, at low-micromolar concentrations, CBD was found to inhibit indoleamine-2,3-dioxygenase activity, thereby suppressing tryptophan degradation by mitogen-stimulated peripheral blood mononuclear cells and LPS-stimulated myelomonocytic THP-1 cells *in vitro* [21]. Based on this finding, CBD might be useful therapeutically to counter the increased risk of depression in diseases associated with

immune activation and inflammation, which often lead to decreased tryptophan, the precursor of serotonin. Finally, CBD has been shown to act as an antagonist at G-protein-coupled receptor 55 and as an antagonist or agonist at several transient receptor potential channels; however, these observations are controversial and the pharmacophysiological significance of these interactions is not known [1,4].

### Actions on immune cells

CBD has been shown to modulate the function of the immune system. Overall these actions may be nuanced and concentration-dependent, but in general include suppression of both cell-mediated and humoral immunity and involve inhibition of proliferation, maturation, and migration of immune cells, antigen presentation, and humoral response [1,13]. Key aspects are discussed here. In most *in vivo* models of inflammation, CBD attenuates inflammatory cell migration/infiltration (e.g., neutrophils) [22]. During neuroinflammation, activated microglial cells migrate toward the site of injury where they release proinflammatory cytokines and cytotoxic agents, including ROS. Although important in the removal of cellular debris and fighting infection, activated microglial cells often exacerbate local cell damage. CBD was shown to inhibit activated microglial cell migration by antagonizing the abnormal-cannabidiol (Abn-CBD)-sensitive receptor at concentrations  $<1 \mu\text{M}$  [23]. Evidence that the Abn-CBD receptor is the orphan G-protein-coupled receptor GPR18 was recently reported [24]. CBD was also shown to block endotoxin-induced oxidative stress resulting from retinal microglial cell activation in uveitis [25]. CBD blocked the immediate activation of NADPH oxidase as well as a second wave of ROS formation and the associated TNF $\alpha$  secretion and p38 MAPK activation. The direct antioxidant property of CBD is unlikely to be the entire explanation for these actions as they occurred at a concentration of  $1 \mu\text{M}$ . Inhibition of adenosine uptake as discussed previously may have been involved. However, a complete understanding of the anti-inflammatory actions of CBD on microglial cells is not yet available. Recently, through an unidentified mechanism, CBD was reported to suppress LPS-induced proinflammatory signaling in cultured microglial cells, including NF- $\kappa\text{B}$  and STAT1 activation, while enhancing STAT3-related anti-inflammatory signaling [26].

CBD induces apoptosis of monocytes and certain normal and transformed lymphocytes, including thymocytes and splenocytes, through oxidative stress and increased ROS levels [27–31]. The basis for this action seems to be glutathione depletion due to adduct formation with the reactive metabolite of CBD, cannabidiol hydroxyquinone, thereby triggering cell death through caspase-8 activation and/or the intrinsic apoptotic pathway. Increased ROS from the upregulation of NADPH oxidases via an undefined mechanism may contribute to cell death as well [31]. A recent study assessed the impact of repeated administration of relatively low levels of CBD to adult male Wistar rats on peripheral blood lymphocyte subset distribution [32]. At 2.5 mg/kg/day for 14 days, CBD did not produce lymphopenia, but increased the total number of natural killer T (NKT) cells and the percentages of NKT and NK cells. A dose of 5 mg/kg/day did have a lymphopenic effect, but by reducing B, T, Tc, and Th lymphocytes. Thus, CBD would seem to suppress specific immunity, while enhancing nonspecific antitumor and antiviral immune response. As discussed by the authors [32], the lymphopenic effect of CBD was observed at a concentration shown to be efficacious in a number of animal models of neurodegenerative and inflammatory diseases, including blocking the progression of collagen-induced arthritis in a murine model of rheumatoid arthritis, decreasing damage to pancreatic islets in the nonobese diabetes-prone (NOD) mouse model of type 1 diabetes, lessening hyperalgesia in rat models of neuropathic and inflammatory pain, and preventing cerebral ischemia in gerbils.

### Pain

Neuropathic pain is associated with microglia activation in the spinal cord and brain and their subsequent release of proinflammatory cytokines, such as interleukin-6 (IL-6), IL-1 $\beta$ , and TNF $\alpha$  [33]. The etiology of neuropathic pain, which is common in cancer, diabetes, multiple sclerosis, and peripheral nerve injury, is poorly understood, but recent evidence indicates that increased ROS generation by microglial cells is the critical initiating factor [34]. The drug Sativex, which consists of  $\Delta^9$ -THC and CBD, is approved in several countries for treatment of central and peripheral neuropathic pain and for spasticity associated with multiple sclerosis [35]. In a mouse model of type 1 diabetic peripheral neuropathic pain, intranasal or intraperitoneal administration of a moderate-high dose of CBD attenuated tactile allodynia and thermal hypersensitivity without affecting the diabetic state [36]. The antinociceptive effects of CBD were associated with less of an increase in microglial density and p38 MAPK activity in the dorsal spinal cord. Finally, the anti-inflammatory and immunosuppressive actions of CBD may be of use in treating rheumatoid arthritis and the associated pain [37,38].

### Diabetes and diabetic complications

CBD was shown to reduce either the initiation of diabetes or the development of overt or latent diabetes in NOD mice by reducing insulinitis [39,40]. This action was accompanied by a shift in the immune response from a dominant Th1 pattern with proinflammatory cytokines to a Th2 pattern with increased levels of the anti-inflammatory cytokine IL-10. Major effectors of  $\beta$ -cell death in type 1 diabetes are various free radicals and oxidant species, including nitric oxide (NO), and infiltrating macrophages are one source of high concentrations of NO and inflammatory cytokines that further enhance NO and ROS formation [41]. CBD was also shown to be effective in blocking ROS-induced upregulation of surface adhesion molecules on endothelial cells due to high glucose and in preserving endothelial barrier function [19,42]. Adhesion of monocytes followed by their transmigration into the subendothelial space is an early event in atherosclerosis, the most common macrovascular complication of diabetes, and may contribute as well to diabetic retinopathy [19,42,43]. The anti-inflammatory actions of CBD may also protect retinal neurons in diabetes by attenuating activation and ROS generation by Müller glia, thus preventing tyrosine nitration and inhibition of Müller cell glutamine synthetase and the consequent accumulation of glutamate, which in turn leads to oxidative stress-induced death of retinal neuronal cells [44].

In a mouse model of type 1 diabetic cardiomyopathy, both pre- and posttreatment with CBD attenuated cardiac fibrosis and cell death, myocardial dysfunction, inflammation, oxidative/nitrosative stress, and the activation of related signaling pathways [45]. CBD attenuated diabetes-induced activation in the heart of the key proinflammatory transcription factor, NF- $\kappa\text{B}$ , and its consequences, e.g., expression of ICAM-1, iNOS, VCAM-1, and TNF $\alpha$ . These observations underscore the point that CBD probably attenuates inflammation far beyond its antioxidant properties *per se*. CBD also reduced high-glucose-induced increases in both cytosolic and mitochondrial reactive oxygen and nitrogen species generation in primary human cardiac myocytes, which was accompanied by reduced NF- $\kappa\text{B}$  activation and cell death. These findings indicate that CBD may have great therapeutic potential in alleviating cardiac complications of diabetes.

### Hypertension

Although CBD has not been considered for treating hypertension, a parallel between the role of microglia in diabetes and hypertension deserves mention. Activation of microglia within the paraventricular nucleus (PVN) was recently shown to contribute to neurogenic

hypertension resulting from chronic angiotensin II infusion in the rat [46]. Microglia activation was associated with enhanced expression of proinflammatory cytokines, the acute administration of which into the left ventricle or PVN resulted in increased blood pressure. The hypertensive action of angiotensin II infusion could be blocked by overexpression of IL-10 in the PVN or intracerebroventricular infusion of minocycline, supporting the involvement of ROS.

The immune system contributes as well to systemic endothelial dysfunction observed in hypertension [47]. Local production of angiotensin II by activated leukocytes within the vessel wall is thought to reduce endothelial function and NO production, leading to attenuated vasodilation and increased blood pressure, through the production of inflammatory cytokines and ROS [48,49]. Interestingly, recent evidence has shown that the initial stimulus for peripheral leukocyte activation in angiotensin II-induced hypertension is the increase in blood pressure that results from stimulation of cells within the anteroventral third ventricle of the brain by angiotensin II [50].

### Ischemia–reperfusion injury

Redox stress and ROS produced by ischemia–reperfusion of organs activates the immune system, which aids in repair by removing debris and stimulating remodeling. An excessive or prolonged inflammatory response, however, may prove detrimental to organ function by exacerbating ROS production and causing death of the parenchyma. Several hours after ischemia–reperfusion in the heart, a model of myocardial infarction, neutrophils accumulate in the myocardium [51]. Several lines of evidence suggest that this accumulation of neutrophils worsens injury to the myocardium [51]. In rats, treatment with CBD for 7 days after a 30-min occlusion of the left anterior descending coronary artery markedly reduced infarct size, myocardial inflammation, and IL-6 levels and preserved cardiac function [52]. In addition, the number of leukocytes infiltrating the border of the infarcted area was dramatically reduced. CBD has been shown to inhibit stimulated migration of neutrophils [22]. CBD treatment was also recently shown to reduce neutrophil migration in a rat model of periodontitis [53]. Hyperactive neutrophils exacerbate periodontal tissue injury and lead to tooth loss in part by excessive ROS formation in individuals with refractory periodontitis [54]. Finally, pre- or postischemic treatment with CBD was shown to have a prolonged and potent protective action in cerebral ischemia. The neuroprotective actions of CBD were attributed to reduced neutrophil accumulation and myeloperoxidase activity [55], as well as decreased high-mobility group box 1 expression by microglia [56].

### Depression

CBD is reported to have antidepressive actions, the basis for which is not established although activation of 5-HT<sub>1A</sub> receptors may be involved at least at higher concentrations [13,57,58]. Growing evidence in recent years has implicated proinflammatory cytokines, free radical species, and oxidants in the etiology of depression [59,60]. One explanation is that the resultant oxidative stress adversely affects glial cell function and leads to neuron damage in the brain.

### Neurodegenerative diseases

Microglial hyperactivation is a common feature of a number of neurodegenerative diseases, including Parkinson, Alzheimer, Huntington, amyotrophic lateral sclerosis, and multiple sclerosis [61,62]. Activated microglia produce a number of pro- and anti-inflammatory cytokines, chemokines, glutamate, neurotrophic factors, and prostanooids and a variety of free radicals that together create a state of oxidative stress. Alzheimer disease, which is the most common form of dementia, is characterized by the deposition of “senile” plaques that are sites of microglia activation and inflammation. The resultant oxidative

stress is a critical factor in the pathophysiology of Alzheimer [63]. The plaques are composed of insoluble aggregates of the  $\beta$ -amyloid peptide (A $\beta$ ), which self-assembles as monomers, oligomers, and finally fibrils. Recent evidence shows that the oligomeric form of  $\beta$ -amyloid is the most neurotoxic species and is most effective as a chemotactic agent for microglia and stimulator of microglial oxidative stress [61,64]. Activated microglia are a major contributor of inflammatory factors in Alzheimer disease and secrete a number of proinflammatory cytokines, which ironically further enhance A $\beta$  production by neuronal cells [65]. In addition, an inflammatory state was shown to block the ability of microglia to phagocytose fibrillar A $\beta$  [66]. Aging was also shown to negatively influence the ability of microglia to internalize A $\beta$  [67]. Microglia from aged mice were also shown to be less responsive to stimulation and to secrete greater amounts of IL-6 and TNF $\alpha$  compared to microglia of younger mice. Aged microglia also had lower levels of glutathione, suggesting an increased susceptibility to the harmful effects of oxidative stress. Finally, although controversial, evidence has been put forward suggesting that bone marrow-derived monocytic cells may somehow gain access to the diseased brain in Alzheimer disease and be better at phagocytosing amyloid plaques than resident microglia [65,68].

Based on rather scant evidence, some have proposed that CBD might have utility in treating neurodegenerative diseases [1,3,69–71]. CBD was shown to have a protective effect on cultured rat pheochromocytoma PC12 cells exposed to A $\beta$  [72,73]. In a concentration-dependent manner, CBD increased cell survival while decreasing ROS and nitrite production, lipid peroxidation, and iNOS protein expression. CBD was shown to have anti-inflammatory actions in vivo in a mouse model of Alzheimer neuroinflammation induced by injection of human A $\beta$  into the hippocampus. CBD dose-dependently attenuated A $\beta$ -induced glial fibrillary acidic protein mRNA, iNOS and IL-1 $\beta$  protein expression, and NO and IL-1 $\beta$  release [74]. In a recent study, CBD was found to protect against amphetamine-induced oxidative protein damage in a rat model of mania and to increase brain-derived neurotrophic factor expression levels in the reversal protocol [75]. Results of these preclinical studies are persuasive and support the need for double-blind placebo-controlled trials to assess the therapeutic utility of CBD in patients with neurodegenerative diseases.

### Obesity and the metabolic syndrome

Metabolic syndrome is a combination of medical disturbances including central obesity, glucose intolerance, hypertension, and dyslipidemia that increases the risk for developing cardiovascular diseases and type 2 diabetes. Adipocyte dysfunction leading to a low-grade chronic inflammatory state is thought to underpin the etiology of the metabolic syndrome [76]. Metabolic overload of adipocytes causes production of ROS, proinflammatory cytokines, and adipokines that activate inflammatory genes and stress kinases and interfere with insulin signaling [76,77]. Saturated fatty acids also activate Toll-like receptors on adipocytes and macrophages, components of the innate immune system, to induce production of proinflammatory cytokines and chemokines. Enhanced mitochondrial flux together with relative hypoxia due to adipocyte tissue hypertrophy, endothelial cell apoptosis, and inflammation-impaired angiogenesis further enhances ROS generation. Enhanced rupture of adipocytes because of excessive hypertrophy attracts and activates macrophages that further exacerbate the inflammatory state through the production of inflammatory cytokines and ROS. The chronic inflammatory state compromises the ability of adipose tissue to absorb incoming fat leading to fat buildup in other organs, including liver, heart, and skeletal muscle, and creating a local inflammatory state that progresses to insulin resistance in those organs as well. Increased ROS levels are thought to be the major contributing factor to insulin resistance [78,79].


Macrophages, both resident and, to a greater extent, bone marrow derived, play a critical role in initiating adipose tissue dysregulation and inflammation in the metabolic syndrome and together with adipocytes constitute a paracrine loop that sustains the chronic inflammatory state [80]. Macrophages secrete TNF $\alpha$ , which acts on hypertrophied adipocytes to downregulate adiponectin and induce proinflammatory cytokines and lipolysis. The released free fatty acids act in turn on the Toll-like receptor 4 of macrophages to induce production of pro-inflammatory cytokines, including TNF $\alpha$ . Both macrophages and adipocytes secrete monocyte chemoattractant protein 1 (MCP1), which serves to recruit more macrophages to the adipose tissue.

Recent evidence has revealed that most macrophages in obese adipose tissue are polarized toward the M1 or classically activated, proinflammatory state, as opposed to the M2 or alternatively activated, anti-inflammatory state [80,81]. The Th1 cytokine interferon- $\gamma$ , microbial by-products (e.g., LPS), and free fatty acids from visceral adipose tissue promote polarization toward the M1 state, whereas the Th2 cytokines IL-4 and IL-13 promote polarization toward the M2 phenotype. The ligand-dependent transcription factors peroxisome proliferator-activated receptors (PPARs) play a key role in determining the M1/M2 phenotype [81,82]. Activation of PPAR $\gamma$  or PPAR $\delta$  promotes differentiation toward the M2 phenotype, and PPAR $\gamma$  activation inhibits the M2 to M1 phenotype switch and represses the M1 proinflammatory gene expression profile. Of interest, CBD, as well as some other cannabinoids, has been shown to activate PPAR $\gamma$ , possibly through direct binding [83,84]. Although tonic activation of CB $_1$  receptors by endocannabinoids is implicated in the development of abdominal obesity, and CB $_1$  antagonists and inverse agonist reduce obesity, their

clinical use is problematic because of serious neuropsychiatric effects [85]. Given its anti-inflammatory actions and PPAR $\gamma$  agonism, CBD might serve as the basis for design of a new antiobesity drug [1]. In this regard, a cautionary note regarding the PPAR $\gamma$  agonism associated with CBD should be sounded, although this was observed only in very high concentrations and only in vitro, which is that several PPAR $\gamma$  agonists have been retracted because of various problems [86,87].

### Atherosclerosis

Atherosclerosis is an inflammatory disease in which monocytes/macrophages play a critical role in the initiation and progression, as well as rupture, of the atherosclerotic plaque [88]. Plaques form in the arterial wall at areas of disturbed flow and endothelial dysfunction (Fig. 2). The initiating event is the transcytosis of low-density lipoprotein (LDL) into the subendothelial space where it is trapped by binding to proteoglycans of the extracellular matrix [88,89]. LDL is oxidized by various cells, including macrophages, first to minimally modified LDL (mmLDL) and then extensively oxidized LDL (oxLDL). The former activates endothelial cells to secrete various factors that attract monocytes and to express adhesion molecules that support the binding and transmigration of monocytes into the subendothelial space. Once there, monocytes differentiate into macrophages under the influence of cytokines and oxLDL. Macrophages take up oxLDL and differentiate into foam cells that secrete a number of cytokines and growth factors that sustain the inflammatory response and stimulate migration of smooth muscle and endothelial cells into the intima. Continued oxLDL uptake by foam cells combined with impaired cholesterol efflux results in their apoptosis and exposure of thrombogenic lipids [88,89]. A number of events in



**Fig. 2.** Inflammation and oxidative stress in atherosclerotic plaque formation. (1) Endothelial dysfunction causes monocyte activation and their binding to endothelial cells, via the production of MCP1, its binding to CCR2 receptors, and the upregulation of adhesion molecules on endothelial cells. (2) Monocytes cross the endothelium and differentiate into macrophages. (3) Because of ROS, LDL that traverses the endothelium is converted to mmLDL and oxLDL. Macrophages accumulate oxLDL through scavenger receptors and are turned into foam cells. (4) Along with T cells, foam cells produce inflammatory mediators that stimulate migration of smooth muscle and endothelial cells into the intima. Reproduced with permission from Ref. [88].

monocyte/macrophage physiology may be potential therapeutic targets for dealing with atherosclerosis and are discussed in detail elsewhere [88,89].

ROS play a pivotal role in atheroma development, and macrophages are the major source for ROS, with NADPH oxidase, cyclooxygenases, lipoxygenases (LOXs), iNOS, and myeloperoxidase contributing [88,89]. ROS participate in atherosclerosis in part by causing LDL oxidation, activating stress signaling pathways, inducing apoptosis, and facilitating plaque rupture [88]. Based on their ability to inhibit 15-LOX, CBD and its mono- and dimethylated derivatives have been proposed as potentially useful in treating atherosclerosis [90]; however, the question of whether 15-LOX has a detrimental or beneficial role in atherosclerosis is unsettled [91]. Nevertheless, a growing body of evidence supports the utility of targeting endocannabinoid signaling, particularly that of macrophages, in the treatment of atherosclerosis [92]. Differentiation of human monocytes, including that induced by oxLDL, results in a change in their CB<sub>1</sub> and CB<sub>2</sub> expression profile such that CB<sub>1</sub> becomes more prominent [93]. Activation of the macrophage CB<sub>1</sub> receptor was shown to upregulate the CD36 scavenger receptor and cholesterol accumulation by macrophages/foam cells [94]. CB<sub>1</sub> receptor activation of human macrophages was linked to ROS generation via p38 MAPK activation, as well as production of TNF $\alpha$  and MCP1 [93]. In contrast, activation of the CB<sub>2</sub> receptor was shown to attenuate the proinflammatory actions of the CB<sub>1</sub> receptor through activation of the Ras-family small G protein Rap1 [93]. Consistent with these findings, a nonselective CB<sub>1</sub>/CB<sub>2</sub> receptor agonist reduced oxLDL-induced ROS generation and TNF $\alpha$  secretion via the CB<sub>2</sub> receptor of murine macrophages, which in contrast to human macrophages do not express much CB<sub>1</sub> receptor [93,95]. Such tantalizing findings have fueled the idea that the endocannabinoid system may be a avenue for further drug development in dealing with atherosclerosis, probably involving a role for CBD as well [7].

Opposing regulatory effects of CB<sub>1</sub> and CB<sub>2</sub> receptors on inflammation and oxidative/nitrative stress are a general theme that has significance in atherosclerosis, as well as other human maladies. CB<sub>2</sub> activation in endothelial cells, which play a key role in development of early atherosclerosis and any inflammatory response, decreases activation and the inflammatory response [96], whereas CB<sub>1</sub> activation in human coronary artery endothelial cells was reported to induce ROS-dependent and -independent MAPK activation and cell death [97]. CB<sub>1</sub> cannabinoid receptors promote oxidative stress and cell death in murine models of doxorubicin-induced cardiomyopathy and in human cardiac myocytes [98]. In contrast, CB<sub>2</sub> activation was found to reduce oxidative stress and neutrophil infiltration in the infarcted mouse myocardium [99]. In nephropathy, CB<sub>2</sub> limits oxidative/nitrosative stress, inflammation, and cell death [100], whereas activation of CB<sub>1</sub> cannabinoid receptors promotes oxidative/nitrosative stress, inflammation, and cell death [101].

## Conclusions

Inflammation and oxidative stress are intimately involved in the genesis of many human diseases. Unraveling that relationship therapeutically has proven challenging, in part because inflammation and oxidative stress “feed off” each other. However, CBD would seem to be a promising starting point for further drug development given its antioxidant (although relatively modest) and anti-inflammatory actions on immune cells, such as macrophages and microglia. CBD also has the advantage of not having psychotropic side effects. Studies on models of human diseases support the idea that CBD attenuates inflammation far beyond its antioxidant properties, for example, by targeting inflammation-related intracellular signaling events. The details on how CBD targets inflammatory signaling remain to be defined. The therapeutic utility of CBD is a relatively new area of investigation that portends new discoveries in the interplay between

inflammation and oxidative stress, a relationship that underlies tissue and organ damage in many human diseases.

## Acknowledgments

This work was supported by grants from the National Heart, Lung, and Blood Institute (R01HL088101-04 and R01HL088101-02S1).

## References

- [1] Izzo, A. A.; Borrelli, F.; Capasso, R.; Di Marzo, V.; Mechoulam, R. Non-psychotropic plant cannabinoids: new therapeutic opportunities from an ancient herb. *Trends Pharmacol. Sci.* **30**:515–527; 2009.
- [2] Mechoulam, R.; Peters, M.; Murillo-Rodriguez, E.; Hanus, L. O. Cannabidiol—recent advances. *Chem. Biodivers.* **4**:1678–1692; 2007.
- [3] Zuardi, A. W. Cannabidiol: from an inactive cannabinoid to a drug with wide spectrum of action. *Rev. Bras. Psiquiatr.* **30**:271–280; 2008.
- [4] De Petrocellis, L.; Di Marzo, V. Non-CB<sub>1</sub>, non-CB<sub>2</sub> receptors for endocannabinoids, plant cannabinoids, and synthetic cannabimimetics: focus on G-protein-coupled receptors and transient receptor potential channels. *J. Neuroimmune Pharmacol.* **5**:103–121; 2010.
- [5] Graham, E. S.; Ashton, J. C.; Glass, M. Cannabinoid receptors: a brief history and “what’s hot”. *Front. Biosci.* **14**:944–957; 2009.
- [6] Tanasescu, R.; Constantinescu, C. S. Cannabinoids and the immune system: an overview. *Immunobiology* **215**:588–597; 2010.
- [7] Pacher, P.; Steffens, S. The emerging role of the endocannabinoid system in cardiovascular disease. *Semin. Immunopathol.* **31**:63–77; 2009.
- [8] Bátkai, S.; Pacher, P. Endocannabinoids and cardiac contractile function: pathophysiological implications. *Pharmacol. Res.* **60**:99–106; 2009.
- [9] Hampson, A. J.; Grimaldi, M.; Lolic, M.; Wink, D.; Rosenthal, R.; Axelrod, J. Neuroprotective antioxidants from marijuana. *Ann. NY Acad. Sci.* **899**:274–282; 2000.
- [10] Chen, Y.; Buck, J. Cannabinoids protect cells from oxidative cell death: a receptor-independent mechanism. *J. Pharmacol. Exp. Ther.* **293**:807–812; 2000.
- [11] Hamelink, C.; Hampson, A.; Wink, D. A.; Eiden, L. E.; Eskay, R. L. Comparison of cannabidiol, antioxidants, and diuretics in reversing binge ethanol-induced neurotoxicity. *J. Pharmacol. Exp. Ther.* **314**:780–788; 2005.
- [12] García-Arencibia, M.; González, S.; de Lago, E.; Ramos, J. A.; Mechoulam, R.; Fernández-Ruiz, J. Evaluation of the neuroprotective effect of cannabinoids in a rat model of Parkinson’s disease: importance of antioxidant and cannabinoid receptor-independent properties. *Brain Res.* **1134**:162–170; 2007.
- [13] Pertwee, R. G. The diverse CB<sub>1</sub> and CB<sub>2</sub> receptor pharmacology of three plant cannabinoids:  $\Delta^9$ -tetrahydrocannabinol, cannabidiol and  $\Delta^9$ -tetrahydrocannabinol. *Br. J. Pharmacol.* **153**:199–215; 2008.
- [14] Capasso, R.; Borrelli, F.; Aviello, G.; Romano, B.; Scalisi, C.; Capasso, F.; Izzo, A. A. Cannabidiol, extracted from *Cannabis sativa*, selectively inhibits inflammatory hypermotility in mice. *Br. J. Pharmacol.* **154**:1001–1008; 2008.
- [15] Liou, G. I.; Auchampach, J. A.; Hillard, C. J.; Zhu, G.; Yousufzai, B.; Mian, S.; Khan, S.; Khalifa, Y. Mediation of cannabidiol anti-inflammation in the retina by equilibrative nucleoside transporter and A<sub>2A</sub> adenosine receptor. *Investig. Ophthalmol. Vis. Sci.* **49**:5526–5531; 2008.
- [16] Carrier, E. J.; Auchampach, J. A.; Hillard, C. J. Inhibition of an equilibrative nucleoside transporter by cannabidiol: a mechanism of cannabinoid immunosuppression. *Proc. Natl Acad. Sci. USA* **103**:7895–7900; 2006.
- [17] Haskó, G.; Csóka, B.; Németh, Z. H.; Vizi, E. S.; Pacher, P. A<sub>2B</sub> adenosine receptors in immunity and inflammation. *Trends Immunol.* **30**:263–270; 2009.
- [18] Haskó, G.; Linden, J.; Cronstein, B.; Pacher, P. Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat. Rev. Drug Discovery* **7**:759–770; 2008.
- [19] Rajesh, M.; Mukhopadhyay, P.; Bátkai, S.; Haskó, G.; Liaudet, L.; Drel, V. R.; Obrosova, I. G.; Pacher, P. Cannabidiol attenuates high glucose-induced endothelial cell inflammatory response and barrier disruption. *Am. J. Physiol. Heart Circ. Physiol.* **293**:H610–H619; 2007.
- [20] Pan, H.; Mukhopadhyay, P.; Rajesh, M.; Patel, V.; Mukhopadhyay, B.; Gao, B.; Haskó, G.; Pacher, P. Cannabidiol attenuates cisplatin-induced nephrotoxicity by decreasing oxidative/nitrosative stress, inflammation, and cell death. *J. Pharmacol. Exp. Ther.* **328**:708–714; 2009.
- [21] Jenny, M.; Santer, E.; Pirich, E.; Schennach, H.; Fuchs, D.  $\Delta^9$ -Tetrahydrocannabinol and cannabidiol modulate mitogen-induced tryptophan degradation and neopterin formation in peripheral blood mononuclear cells *in vitro*. *J. Neuroimmunol.* **207**:75–82; 2009.
- [22] McHugh, D.; Tanner, C.; Mechoulam, R.; Pertwee, R. G.; Ross, R. A. Inhibition of human neutrophil chemotaxis by endogenous cannabinoids and phytocannabinoids: evidence for a site distinct from CB<sub>1</sub> and CB<sub>2</sub>. *Mol. Pharmacol.* **73**:441–450; 2008.
- [23] Walter, L.; Franklin, A.; Witting, A.; Wade, C.; Xie, Y.; Kunos, G.; Mackie, K.; Stella, N. Nonpsychotropic cannabinoid receptors regulate microglial cell migration. *J. Neurosci.* **23**:1398–1405; 2003.
- [24] McHugh, D.; Hu, S. S.; Rimmerman, N.; Juknat, A.; Vogel, Z.; Walker, J. M.; Bradshaw, H. B. N-arachidonoyl glycine, an abundant endogenous lipid, potently drives directed cellular migration through GPR18, the putative abnormal cannabidiol receptor. *BMC Neurosci.* **26**:44; 2010.

- [25] El-Remessy, A. B.; Tang, Y.; Zhu, G.; Matragoon, S.; Khalifa, Y.; Liu, E. K.; Liu, J. Y.; Hanson, E.; Mian, S.; Fatteh, N.; Liou, G. I. Neuroprotective effects of cannabidiol in endotoxin-induced uveitis: critical role of p38 MAPK activation. *Mol. Vis.* **14**: 2190–2203; 2008.
- [26] Kozela, E.; Pietri, M.; Juknat, A.; Rimmerman, N.; Levy, R.; Vogel, Z. Cannabinoids  $\Delta^9$ -tetrahydrocannabinol and cannabidiol differentially inhibit the lipopolysaccharide-activated NF- $\kappa$ B and interferon- $\beta$ /STAT proinflammatory pathways in BV-2 microglial cells. *J. Biol. Chem.* **285**:1616–1626; 2010.
- [27] Wu, H. Y.; Chang, A. C.; Wang, C. C.; Kuo, F. H.; Lee, C. Y.; Liu, D. Z.; Jan, T. R. Cannabidiol induced a contrasting pro-apoptotic effect between freshly isolated and precultured human monocytes. *Toxicol. Appl. Pharmacol.* **246**:141–147; 2011.
- [28] Wu, H. Y.; Chu, R. M.; Wang, C. C.; Lee, C. Y.; Lin, S. H.; Jan, T. R. Cannabidiol-induced apoptosis in primary lymphocytes is associated with oxidative stress-dependent activation of caspase-8. *Toxicol. Appl. Pharmacol.* **226**:260–270; 2008.
- [29] Lee, C. Y.; Wey, S. P.; Liao, M. H.; Hsu, W. L.; Wu, H. Y.; Jan, T. R. A comparative study on cannabidiol-induced apoptosis in murine thymocytes and EL-4 thymoma cells. *Int. Immunopharmacol.* **8**:732–740; 2008.
- [30] Wu, H. Y.; Jan, T. R. Cannabidiol hydroxyquinone-induced apoptosis of splenocytes is mediated predominantly by thiol depletion. *Toxicol. Lett.* **195**: 68–74; 2010.
- [31] McCallip, R. J.; Jia, W.; Schlomer, J.; Warren, J. W.; Nagarkatti, P. S.; Nagarkatti, M. Cannabidiol-induced apoptosis in human leukemia cells: a novel role of cannabidiol in the regulation of p22<sup>phox</sup> and Nox4 expression. *Mol. Pharmacol.* **70**:897–908; 2006.
- [32] Ignatowska-Jankowska, B.; Jankowski, M.; Glac, W.; Swiergel, A. H. Cannabidiol-induced lymphopenia does not involve NKT and NK cells. *J. Physiol. Pharmacol.* **60** (Suppl. 3):99–103; 2009.
- [33] Baron, R.; Binder, A.; Wasner, G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol.* **9**:807–819; 2010.
- [34] Kim, D.; You, B.; Jo, E. K.; Han, S. K.; Simon, M. I.; Lee, S. J. NADPH oxidase 2-derived reactive oxygen species in spinal cord microglia contribute to peripheral nerve injury-induced neuropathic pain. *Proc. Natl. Acad. Sci. USA* **107**: 14851–14856; 2010.
- [35] Rahn, E. J.; Hohmann, A. G. Cannabinoids as pharmacotherapies for neuropathic pain: from the bench to the bedside. *Neurotherapeutics* **6**:713–737; 2009.
- [36] Toth, C. C.; Jedrzejewski, N. M.; Ellis, C. L.; Frey, W. H. 2nd. Cannabinoid-mediated modulation of neuropathic pain and microglial accumulation in a model of murine type I diabetic peripheral neuropathic pain. *Mol. Pain* **6**:16; 2010.
- [37] Blake, D. R.; Robson, P.; Ho, M.; Jubb, R. W.; McCabe, C. S. Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford)* **45**: 50–52; 2006.
- [38] Malfait, A. M.; Gallily, R.; Sumariwalla, P. F.; Malik, A. S.; Andreaskos, E.; Mechoulam, R.; Feldmann, M. The nonpsychoactive cannabis constituent cannabidiol is an oral anti-arthritis therapeutic in murine collagen-induced arthritis. *Proc. Natl. Acad. Sci. USA* **97**:9561–9566; 2000.
- [39] Weiss, L.; Zeira, M.; Reich, S.; Har-Noy, M.; Mechoulam, R.; Slavin, S.; Gallily, R. Cannabidiol lowers incidence of diabetes in non-obese diabetic mice. *Autoimmunity* **39**:143–151; 2006.
- [40] Weiss, L.; Zeira, M.; Reich, S.; Slavin, S.; Raz, I.; Mechoulam, R.; Gallily, R. Cannabidiol arrests onset of autoimmune diabetes in NOD mice. *Neuropharmacology* **54**:244–249; 2008.
- [41] Pacher, P.; Beckman, J. S.; Liaudet, L. Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* **87**:315–342; 2007.
- [42] El-Remessy, A. B.; Al-Shabraway, M.; Khalifa, Y.; Tsai, N. T.; Caldwell, R. B.; Liou, G. I. Neuroprotective and blood–retinal barrier-preserving effects of cannabidiol in experimental diabetes. *Am. J. Pathol.* **168**:235–244; 2006.
- [43] Noda, K.; Nakao, S.; Zandi, S.; Engelstädter, V.; Mashima, Y.; Hafezi-Moghadam, A. Vascular adhesion protein-1 regulates leukocyte transmigration rate in the retina during diabetes. *Exp. Eye Res.* **89**:774–781; 2009.
- [44] El-Remessy, A. B.; Khalifa, Y.; Ola, S.; Ibrahim, A. S.; Liou, G. I. Cannabidiol protects retinal neurons by preserving glutamine synthetase activity in diabetes. *Mol. Vis.* **16**:1487–1495; 2010.
- [45] Rajesh, M.; Mukhopadhyay, P.; Båtkai, S.; Patel, V.; Saito, K.; Matsumoto, S.; Kashiwaya, Y.; Horváth, B.; Mukhopadhyay, B.; Becker, L.; Haskó, G.; Liaudet, L.; Wink, D. A.; Veves, A.; Mechoulam, R.; Pacher, P. Cannabidiol attenuates cardiac dysfunction, oxidative stress, fibrosis, and inflammatory and cell death signaling pathways in diabetic cardiomyopathy. *J. Am. Coll. Cardiol.* **56**:2115–2125; 2010.
- [46] Shi, P.; Diez-Freire, C.; Jun, J. Y.; Qi, Y.; Katovich, M. J.; Li, Q.; Sriramula, S.; Francis, J.; Summers, C.; Raizada, M. K. Brain microglial cytokines in neurogenic hypertension. *Hypertension* **56**:297–303; 2010.
- [47] Harrison, D. G.; Vinh, A.; Lob, H.; Madhur, M. S. Role of the adaptive immune system in hypertension. *Curr. Opin. Pharmacol.* **10**:203–207; 2010.
- [48] De Miguel, C.; Das, S.; Lund, H.; Mattson, D. L. T lymphocytes mediate hypertension and kidney damage in Dahl salt-sensitive rats. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **298**:R1136–R1142; 2010.
- [49] Azekoshi, Y.; Yasu, T.; Watanabe, S.; Tagawa, T.; Abe, S.; Yamakawa, K.; Uehara, Y.; Momomura, S.; Urata, H.; Ueda, S. Free fatty acid causes leukocyte activation and resultant endothelial dysfunction through enhanced angiotensin II production in mononuclear and polymorphonuclear cells. *Hypertension* **56**:136–142; 2010.
- [50] Marvar, P. J.; Thabet, S. R.; Guzik, T. J.; Lob, H. E.; McCann, L. A.; Weyand, C.; Gordon, F. J.; Harrison, D. G. Central and peripheral mechanisms of T-lymphocyte activation and vascular inflammation produced by angiotensin II-induced hypertension. *Circ. Res.* **107**:263–270; 2010.
- [51] Chao, W. Toll-like receptor signaling: a critical modulator of cell survival and ischemic injury in the heart. *Am. J. Physiol. Heart Circ. Physiol.* **296**:H1–H12; 2009.
- [52] Durst, R.; Danenberg, H.; Gallily, R.; Mechoulam, R.; Meir, K.; Grad, E.; Beeri, R.; Pugatsch, T.; Tarsish, E.; Lotan, C. Cannabidiol, a nonpsychoactive *Cannabis* constituent, protects against myocardial ischemic reperfusion injury. *Am. J. Physiol. Heart Circ. Physiol.* **293**:H3602–H3607; 2007.
- [53] Napimoga, M. H.; Benatti, B. B.; Lima, F. O.; Alves, P. M.; Campos, A. C.; Pena-Dos-Santos, D. R.; Severino, F. P.; Cunha, F. Q.; Guimarães, F. S. Cannabidiol decreases bone resorption by inhibiting RANK/RANKL expression and pro-inflammatory cytokines during experimental periodontitis in rats. *Int. Immunopharmacol.* **9**: 216–222; 2009.
- [54] Johnstone, A. M.; Koh, A.; Goldberg, M. B.; Glogauer, M. A hyperactive neutrophil phenotype in patients with refractory periodontitis. *J. Periodontol.* **78**: 1788–1794; 2007.
- [55] Hayakawa, K.; Mishima, K.; Nozako, M.; Hazekawa, M.; Irie, K.; Fujioka, M.; Orito, K.; Abe, K.; Hasebe, N.; Egashira, N.; Iwasaki, K.; Fujiwara, M. Delayed treatment with cannabidiol has a cerebroprotective action via a cannabinoid receptor-independent myeloperoxidase-inhibiting mechanism. *J. Neurochem.* **102**: 1488–1496; 2007.
- [56] Hayakawa, K.; Irie, K.; Sano, K.; Watanabe, T.; Higuchi, S.; Enoki, M.; Nakano, T.; Harada, K.; Ishikane, S.; Ikeda, T.; Fujioka, M.; Orito, K.; Iwasaki, K.; Mishima, K.; Fujiwara, M. Therapeutic time window of cannabidiol treatment on delayed ischemic damage via high-mobility group box 1-inhibiting mechanism. *Biol. Pharm. Bull.* **32**:1538–1544; 2009.
- [57] El-Alfy, A. T.; Ivey, K.; Robinson, K.; Ahmed, S.; Radwan, M.; Slade, D.; Khan, I.; ElSohly, M.; Ross, S. Antidepressant-like effect of  $\Delta^9$ -tetrahydrocannabinol and other cannabinoids isolated from *Cannabis sativa* L. *Pharmacol. Biochem. Behav.* **95**:434–442; 2010.
- [58] Zanelati, T. V.; Biojone, C.; Moreira, F. A.; Guimarães, F. S.; Joca, S. R. Antidepressant-like effects of cannabidiol in mice: possible involvement of 5-HT<sub>1A</sub> receptors. *Br. J. Pharmacol.* **159**:122–128; 2010.
- [59] Maes, M.; Galecki, P.; Chang, Y. S.; Berk, M. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **35**:676–692; 2011.
- [60] Song, C.; Wang, H. Cytokines mediated inflammation and decreased neurogenesis in animal models of depression. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **35**:760–768; 2011.
- [61] Heurtaux, T.; Michelucci, A.; Losciuto, S.; Gallotti, C.; Felten, P.; Dorban, G.; Grandbarbe, L.; Morga, E.; Heuschling, P. Microglial activation depends on beta-amyloid conformation: role of the formylpeptide receptor 2. *J. Neurochem.* **114**: 576–586; 2010.
- [62] Heneka, M. T.; Rodríguez, J. J.; Verkhratsky, A. Neuroglia in neurodegeneration. *Brain Res. Rev.* **63**:189–211; 2010.
- [63] Candore, G.; Bulati, M.; Caruso, C.; Castiglia, L.; Colonna-Romano, G.; Di Bona, D.; Duro, G.; Lio, D.; Matranga, D.; Pellicanò, M.; Rizzo, C.; Scapagnini, G.; Vasto, S. Inflammation, cytokines, immune response, apolipoprotein E, cholesterol, and oxidative stress in Alzheimer disease: therapeutic implications. *Rejuvenation Res.* **13**:301–313; 2010.
- [64] Tamagno, E.; Bardini, P.; Guglielmo, M.; Danni, O.; Tabaton, M. The various aggregation states of  $\beta$ -amyloid 1–42 mediate different effects on oxidative stress, neurodegeneration, and BACE-1 expression. *Free Radic. Biol. Med.* **41**: 202–212; 2006.
- [65] Sastre, M.; Klockgether, T.; Heneka, M. T. Contribution of inflammatory processes to Alzheimer's disease: molecular mechanisms. *Int. J. Dev. Neurosci.* **24**:167–176; 2006.
- [66] Lee, C. Y.; Landreth, G. E. The role of microglia in amyloid clearance from the AD brain. *J. Neural Transm.* **117**:949–960; 2010.
- [67] Njie, E. G.; Boelen, E.; Stassen, F. R.; Steinbusch, H. W.; Borchelt, D. R.; Streit, W. J. Ex vivo cultures of microglia from young and aged rodent brain reveal age-related changes in microglial function. *Neurobiol. Aging* (in press).
- [68] Malm, T.; Koistinaho, M.; Muona, A.; Magga, J.; Koistinaho, J. The role and therapeutic potential of monocytic cells in Alzheimer's disease. *Glia* **58**:889–900; 2010.
- [69] Bisogno, T.; Di Marzo, V. The role of the endocannabinoid system in Alzheimer's disease: facts and hypotheses. *Curr. Pharm. Des.* **14**:2299–3305; 2008.
- [70] Zuardi, A. W.; Crippa, J. A.; Hallak, J. E.; Pinto, J. P.; Chagas, M. H.; Rodrigues, G. G.; Dursun, S. M.; Tumas, V. Cannabidiol for the treatment of psychosis in Parkinson's disease. *J. Psychopharmacol.* **23**:979–983; 2009.
- [71] Iuvone, T.; Esposito, G.; De Filippis, D.; Scuderi, C.; Steardo, L. Cannabidiol: a promising drug for neurodegenerative disorders? *CNS Neurosci. Ther.* **15**:65–75; 2009.
- [72] Iuvone, T.; Esposito, G.; Esposito, R.; Santamaria, R.; Di Rosa, M.; Izzo, A. A. Neuroprotective effect of cannabidiol, a non-psychoactive component from *Cannabis sativa*, on  $\beta$ -amyloid-induced toxicity in PC12 cells. *J. Neurochem.* **89**: 134–141; 2004.
- [73] Esposito, G.; De Filippis, D.; Maiuri, M. C.; De Stefano, D.; Carnuccio, R.; Iuvone, T. Cannabidiol inhibits inducible nitric oxide synthase protein expression and nitric oxide production in beta-amyloid stimulated PC12 neurons through p38 MAP kinase and NF- $\kappa$ B involvement. *Neurosci. Lett.* **399**:91–95; 2006.
- [74] Esposito, G.; Scuderi, C.; Savani, C.; Steardo Jr., L.; De Filippis, D.; Cottone, P.; Iuvone, T.; Cuomo, V.; Steardo, L. Cannabidiol in vivo blunts  $\beta$ -amyloid induced neuroinflammation by suppressing IL-1 $\beta$  and iNOS expression. *Br. J. Pharmacol.* **151**:1272–1279; 2007.
- [75] Valvassori, S.; Elias, G.; de Souza, B.; Petronilho, F.; Dal-Pizzol, F.; Kapczinski, F.; Trzesniak, C.; Tumas, V.; Dursun, S.; Chagas, M. H.; Hallak, J. E.; Zuardi, A. W.; Quevedo, J.; Crippa, J. A. Effects of cannabidiol on amphetamine-induced oxidative stress generation in an animal model of mania. *J. Psychopharmacol.* **25**:274–280; 2011.
- [76] Monteiro, R.; Azevedo, I. Chronic inflammation in obesity and the metabolic syndrome. *Mediat. Inflamm.* **2010**:289645; 2010.

- [77] Karalis, K. P.; Giannogonas, P.; Kodela, E.; Koutmani, Y.; Zoumakis, M.; Teli, T. Mechanisms of obesity and related pathology: linking immune responses to metabolic stress. *FEBS J.* **276**:5747–5754; 2009.
- [78] Houstis, N.; Rosen, E. D.; Lander, E. S. Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature* **440**:944–948; 2006.
- [79] Samocha-Bonet, D.; Heilbronn, L. K.; Lichtenberg, D.; Campbell, L. V. Does skeletal muscle oxidative stress initiate insulin resistance in genetically predisposed individuals? *Trends Endocrinol. Metab.* **21**:83–88; 2010.
- [80] Suganami, T.; Ogawa, Y. Adipose tissue macrophages: their role in adipose tissue remodeling. *J. Leukoc. Biol.* **88**:33–39; 2010.
- [81] Fuentes, L.; Roszer, T.; Ricote, M. Inflammatory mediators and insulin resistance in obesity: role of nuclear receptor signaling in macrophages. *Mediat. Inflamm.* **2010**:219583; 2010.
- [82] Chawla, A. Control of macrophage activation and function by PPARs. *Circ. Res.* **106**:1559–1569; 2010.
- [83] O'Sullivan, S. E.; Kendall, D. A. Cannabinoid activation of peroxisome proliferator-activated receptors: potential for modulation of inflammatory disease. *Immunobiology* **215**:611–616; 2010.
- [84] O'Sullivan, S. E.; Sun, Y.; Bennett, A. J.; Randall, M. D.; Kendall, D. A. Time-dependent vascular actions of cannabidiol in the rat aorta. *Eur. J. Pharmacol.* **612**:61–68; 2009.
- [85] Bermudez-Silva, F. J.; Viveros, M. P.; McPartland, J. M.; Rodriguez de Fonseca, F. The endocannabinoid system, eating behavior and energy homeostasis: the end or a new beginning? *Pharmacol. Biochem. Behav.* **95**:375–382; 2010.
- [86] Kaul, S.; Bolger, A. F.; Herrington, D.; Giugliano, R. P.; Eckel, R. H. Thiazolidinedione drugs and cardiovascular risks: a science advisory from the American Heart Association and American College of Cardiology Foundation. *Circulation* **121**:1868–1877; 2010.
- [87] Riche, D. M.; Travis King, S. Bone loss and fracture risk associated with thiazolidinedione therapy. *Pharmacotherapy* **30**:716–727; 2010.
- [88] Saha, P.; Modarai, B.; Humphries, J.; Mattock, K.; Waltham, M.; Burnand, K. G.; Smith, A. The monocyte/macrophage as a therapeutic target in atherosclerosis. *Curr. Opin. Pharmacol.* **9**:109–118; 2009.
- [89] Hulsmans, M.; Holvoet, P. The vicious circle between oxidative stress and inflammation in atherosclerosis. *J. Cell. Mol. Med.* **14**:70–78; 2010.
- [90] Takeda, S.; Usami, N.; Yamamoto, I.; Watanabe, K. Cannabidiol-2', 6'-dimethyl ether, a cannabidiol derivative, is a highly potent and selective 15-lipoxygenase inhibitor. *Drug Metab. Dispos.* **37**:1733–1737; 2009.
- [91] Hersberger, M. Potential role of the lipoxygenase derived lipid mediators in atherosclerosis: leukotrienes, lipoxins and resolvins. *Clin. Chem. Lab. Med.* **48**:1063–1073; 2010.
- [92] Immenschuh, S. Endocannabinoid signalling as an anti-inflammatory therapeutic target in atherosclerosis: does it work? *Cardiovasc. Res.* **84**:341–342; 2009.
- [93] Han, K. H.; Lim, S.; Ryu, J.; Lee, C. W.; Kim, Y.; Kang, J. H.; Kang, S. S.; Ahn, Y. K.; Park, C. S.; Kim, J. J. CB1 and CB2 cannabinoid receptors differentially regulate the production of reactive oxygen species by macrophages. *Cardiovasc. Res.* **84**:378–386; 2009.
- [94] Jiang, L. S.; Pu, J.; Han, Z. H.; Hu, L. H.; He, B. Role of activated endocannabinoid system in regulation of cellular cholesterol metabolism in macrophages. *Cardiovasc. Res.* **81**:805–813; 2009.
- [95] Hao, M. X.; Jiang, L. S.; Fang, N. Y.; Pu, J.; Hu, L. H.; Shen, L. H.; Song, W.; He, B. The cannabinoid WIN55, 212–2 protects against oxidized LDL-induced inflammatory response in murine macrophages. *J. Lipid Res.* **51**:2181–2190; 2010.
- [96] Rajesh, M.; Mukhopadhyay, P.; Bátkai, S.; Haskó, G.; Liaudet, L.; Huffman, J. W.; Csiszar, A.; Ungvari, Z.; Mackie, K.; Chatterjee, S.; Pacher, P. CB<sub>2</sub>-receptor stimulation attenuates TNF- $\alpha$ -induced human endothelial cell activation, transendothelial migration of monocytes, and monocyte–endothelial adhesion. *Am. J. Physiol. Heart Circ. Physiol.* **293**:H2210–H2218; 2007.
- [97] Rajesh, M.; Mukhopadhyay, P.; Haskó, G.; Liaudet, L.; Mackie, K.; Pacher, P. Cannabinoid-1 receptor activation induces reactive oxygen species-dependent and -independent mitogen-activated protein kinase activation and cell death in human coronary artery endothelial cells. *Br. J. Pharmacol.* **160**:688–700; 2010.
- [98] Mukhopadhyay, P.; Rajesh, M.; Bátkai, S.; Patel, V.; Kashiwaya, Y.; Liaudet, L.; Evgenov, O. V.; Mackie, K.; Haskó, G.; Pacher, P. CB<sub>1</sub> cannabinoid receptors promote oxidative stress and cell death in murine models of doxorubicin-induced cardiomyopathy and in human cardiomyocytes. *Cardiovasc. Res.* **85**:773–784; 2010.
- [99] Montecucco, F.; Lenglet, S.; Braunersreuther, V.; Burger, F.; Pelli, G.; Bertolotto, M.; Mach, F.; Steffens, S. CB<sub>2</sub> cannabinoid receptor activation is cardioprotective in a mouse model of ischemia/reperfusion. *J. Mol. Cell. Cardiol.* **46**:612–620; 2009.
- [100] Mukhopadhyay, P.; Rajesh, M.; Pan, H.; Patel, V.; Mukhopadhyay, B.; Bátkai, S.; Gao, B.; Haskó, G.; Pacher, P. Cannabinoid-2 receptor limits inflammation, oxidative/nitrosative stress, and cell death in nephropathy. *Free Radic. Biol. Med.* **48**:457–467; 2010.
- [101] Mukhopadhyay, P.; Pan, H.; Rajesh, M.; Bátkai, S.; Patel, V.; Harvey-White, J.; Mukhopadhyay, B.; Haskó, G.; Gao, B.; Mackie, K.; Pacher, P. CB<sub>1</sub> cannabinoid receptors promote oxidative/nitrosative stress, inflammation and cell death in a murine nephropathy model. *Br. J. Pharmacol.* **160**:657–668; 2010.